

CLTC[®] 100 MR

Type A Medicated Article

Description:

CLTC 100MR Type A Medicated Article is a dispersible formulation of Chlortetracycline for use in calves. It is a broad-spectrum anti-infective that is effective against a wide variety of infectious diseases caused by susceptible Gram-positive and Gram-negative bacteria. It is safe, stable and highly effective when used as directed.

Active Ingredient:

Chlortetracycline hydrochloride100 g/lb

Mixing and Use Directions:

For premix quantities less than 1 lb. mix the proper amount of premix with at least 10 lb of feed ingredients to make a preblend. Add the preblend to the remaining feed ingredients and mix thoroughly. See table 1 for registered claim and usage directions for **CLTC 100 MR**.

For use in dry feeds only. Not for use in liquid feed supplements.

Store at or below 25°C (77°F), Excursions permitted up to 40°C (104°F)

Caution:

Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated animal feeds only.

Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions - seek prompt medical treatment if such reactions are suspected.

Table 1. Indications for Use	Use Level of Chlortetracycline hydrochloride	lb of CLTC 100 MR /ton
CALVES (milk replacer or starter feeds)		
For calves (up to 250 lb) for treatment of bacterial enteritis caused by <i>E. coli</i> susceptible to chlortetracycline.	10 mg/lb of body weight daily Limit use to 5 days	20 ¹
<p>Warning: At 10 mg/lb use level, withdraw 10 days before slaughter. A withdrawal period has not been established in preruminating calves. Do not use in calves to be processed for veal.</p>		
<p>¹ Calf weighing 100 lb consuming 1 lb of dry starter feed or 1 lb of milk replacer mixed with one gallon of water.</p>		

Not for human use.

Restricted drug (California) - Use only as directed.

HEALTHY ANIMALS. HEALTHY FOOD. HEALTHY WORLD.®

