

Coxistac® 12% Granular

Medicated Feed Article

(DIN# 02237631)

Coxistac 12% Granular is an anticoccidial medication for broiler feeds, containing the polyether ionophore, salinomycin sodium.

Coxistac enables broilers to reach market weight faster with reduced losses from coccidiosis. This means more efficient and profitable production.

Active Ingredient:

Salinomycin sodium..... 12%
(Contains 120 g/kg)

Description:

Off-white to beige granular particle.

Formulation:

Coxistac is a 12% granulated form of salinomycin sodium.

Classification:

Coxistac (salinomycin sodium) 12% Granular is listed in the Medicating Ingredients Brochure (M.I.B.):
Broilers: M.I.B. No. 69

Features:

- Broad anticoccidial coverage against the six major *Eimeria* species, including *E. acervulina*, *E. tenella*, *E. maxima*, *E. brunetti*, *E. necatrix*, *E. mitis* in broiler chickens. Coccidia are destroyed rather than inhibited.
- Acts early in the coccidial life cycle: it destroys sporozoites, trophozoites and early developing schizonts before birds suffer damage.
- Excellent "efficacy margin": the recommended dose level provides a high level of protection through temporary periods of low feed intake caused by disease or climate.
- No withdrawal period
- Good feed mixability

Mixing Directions:

Coxistac medication must be thoroughly mixed in the feed to assure even distribution.

Thoroughly mix 0.5 kg of **Coxistac** 12% Granular with 1,000 kg of complete feed, to obtain a medicated feed with salinomycin sodium concentration of 0.006% (60 mg/kg). The medicated feed can be prepared in the form of meal or pellets.

Safety:

Coxistac medicated feeds, when used as directed, can be safely administered to broiler chickens without fear of animal toxicity, residue problems or environmental pollution.

Care should be taken to avoid contamination of feed for turkeys, dogs and horses. This drug is known to be toxic to these species.

Do not administer tiamulin to broilers receiving feeds medicated with **Coxistac**.

Warning:

Do not feed to replacement, laying or breeding chickens.

Claims and Use Directions

Indication for Use in Broiler Chickens	Dosage and Administration	Directions
As an aid in the prevention of coccidiosis in broiler chickens	Level of drug: 60 ppm (60 mg/kg) of complete feed	Feed to broiler chickens. No withdrawal period required.

Solutions totales



Total Solutions

HEALTHY ANIMALS. HEALTHY FOOD. HEALTHY WORLD.®



Distributed by Bio Agri Mix LP, 11 Ellen Street, Michell Ontario, N0K 1N0
Order Desk: 1.800.265.1763 • Order Fax: 1.519.348.4100 • www.phibroah.com

© 2017 Phibro Animal Health Corporation

Coxistac, Phibro, Phibro logo design and Healthy Animals. Healthy Food. Healthy World are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

CX17001/CAN/0217