



## **Terramycin** (oxytetracycline) Veterinary Feed Directive for use in lobster

Client:	Veterinarian:
Business or Home Address:	Address:
Phone #:	Phone #:
Approximate number of <i>lobsters</i> to be treated: Locations of animals:	
Special instructions and/or other animal identifications:	
Indications:	
For the use in lobsters for the control of gaffkemia caused by <i>Aero</i>	coccus viridans
To the use in leading to the use serials of gamerina saucea by here	seessa viinaane.
Drug Level in Medicated Feed and Duration of Use:	
Level of Drug: 1g/lb	
Duration of Use: 5 consecutive days	
Use of feed containing this veterinary feed directive (VFD) drug in a manner other than as directed on the labeling	
(extralabel use) is not permitted.	
Davidus Wassians Wildedness and diseased	£
Residue Warning: Withdraw medicated	feed 30 days before harvesting lobsters.
Combination Use: This VED only authorizes the use of the VED drug(s) cited in this of	order and is not intended to authorize the use of such drug(s) in
This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.	
VFD Issue Date:	VFD Expiration Date:
Veterinarian's Signature:	month/day/year
	(Not to exceed 6 months from issuance date)

NADA 038-439 Approved by FDA 038-439-VFD-ALB-01 Revised Sept. 20, 2016