



# **Neo-Terramycin** (neomycin-oxytetracycline) Veterinary Feed Directive for use in calves, beef cattle, and poplectating dainy cattle

Client: Business or Home Address:		Veterinarian: Address:	
Phone #:		Phone #:	
Approximate number of <i>animals</i> to Locations of animals:			
Special instructions and/or other a	nimal identifications:		

#### Indications:

- **A.)** For growing cattle (over 400 lb) for reduction of liver condemnation due to liver abscesses
- **B.)** For cattle for the prevention and treatment of the early stages of shipping fever complex.
- C.) For calves and beef and nonlactating dairy cattle for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida* susceptible to oxytetracycline: treatment and control of colibacillosis (bacterial enteritis) caused by *E. coli* susceptible to neomycin.
- **D.)** For calves (up to 250 lb) for the treatment of bacterial enteritis caused by *E. coli* susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by *E. coli* susceptible to neomycin.

### Drug Level in Medicated Feed and Duration of Use:

- A.) Level of Drug: \_\_\_\_\_\_g/ton (specify level to provide 75 mg/head/day)
- Duration of Use: \_\_\_\_\_days (specify number)
- **B.)** Level of Drug: \_\_\_\_\_\_g/ton (specify level to provide 0.5-2.0 g/head/day)
- Duration of Use: \_\_\_\_\_\_days (specify number between 3-5 days before and after arrival in feedlots)
- C.) Level of Drug (in feed or milk replacer): \_\_\_\_\_g/ton (specify level to provide 10 mg/lb BW/day) Duration of Use: \_\_\_\_\_days (specify number between 7-14 days)
- D.) Level of Drug (in feed or milk replacer): \_\_\_\_\_\_g/ton (specify level to provide 10 mg/lb BW/day)

Duration of Use: \_\_\_\_\_\_days (specify number between 7-14 days)

For use in dry feeds only. Not for use in liquid feed supplements. Not for human use. Restricted drug (California) - use only as directed.

Use of feed containing this veterinary feed directive (VFD) drug in a manner other than as directed on the labeling (extralabel use) is not permitted.

## **Residue Warning:**

For indication A there is a zero-day withdrawal period. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

For indication B there is a zero- day withdrawal period. A withdrawal period has not been established in preruminating calves; do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle; do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

For indications C and D withdraw 5 days before slaughter. A withdrawal period has not been established in preruminating calves; do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle; do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

### **Combination Use:**

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

VFD	Issue	Date

Veterinarian's Signature:

NADA 094-975 Approved by FDA

094-975-VFD-CTL-01 Revised Sept. 19, 2016 VFD Expiration Date:

(Not to exceed 6 months from issuance date)

Copy - Client

Phibro Animal Health Corporation 300 Frank W. Burr Blvd. Teaneck, NJ 07666