



Neo-Terramycin (*neomycin-oxytetracycline*) Veterinary Feed Directive for use in turkeys

Client: _____ Veterinarian: _____
 Business or _____ Address: _____
 Home Address: _____
 Phone #: _____ Phone #: _____

Approximate number of turkeys to be treated: _____
 Locations of animals: _____

Special instructions and/or other animal identifications:

Indications:

- A.)** For turkeys for the control of hexamitiasis caused by *Hexamita meleagridis* susceptible to oxytetracycline.
- B.)** For turkeys for the control of infectious synovitis caused by *Mycoplasma synovia* susceptible to oxytetracycline
- C.)** For turkeys for the control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.

Drug Level in Medicated Feed and Duration of Use:

- A.)** Level of Drug: 100 g/ton
- B.)** Level of Drug: 200 g/ton
- C.)** Level of Drug: _____g/ton (specify level to provide 25 mg/lb of BW/day)

Duration of Use: _____days (specify number between 7- 14 days)

For use in dry feeds only. Not for use in liquid feed supplements. Not for human use. Restricted drug (California) - use only as directed.

Use of feed containing this veterinary feed directive (VFD) drug in a manner other than as directed on the labeling (extralabel use) is not permitted.

Residue Warning:

For indication A there is a zero-day withdrawal period. Do not feed to turkeys producing eggs for human consumption.

For indications B and C withdraw 5 days before slaughter. Do not feed to turkeys producing eggs for human consumption.

Combination Use:

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

VFD Issue Date: _____

VFD Expiration Date: _____

Veterinarian's Signature: _____

month/day/year

(Not to exceed 6 months from issuance date)