



Terramycin (oxytetracycline) Veterinary Feed Directive for use in turkeys

Client: _____ Veterinarian: _____
 Business or _____ Address: _____
 Home Address: _____
 Phone #: _____ Phone #: _____

Approximate number of turkeys to be treated: _____
 Location of animals: _____

Special instructions and/or other animal identifications:

Indications, Drug Level and Duration of Use (select one and specify additional required information):

- A.)** For use in turkeys for the control of hexamitiasis caused by *Hexamita meleagridis* susceptible to oxytetracycline
 Level of Drug: 100 g/ton
 Duration of Use: _____ days (7-14 days)
- B.)** For use in turkeys for the control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to oxytetracycline
 Level of Drug: 200 g/ton
 Duration of Use: _____ days (7-14 days)
- C.)** For use in turkeys for the control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline
 Level of Drug: _____g/ton (to provide 25 mg/lb of BW/day)
 Duration of Use: _____ days (7- 14 days)

Use of feed containing this veterinary feed directive (VFD) drug in a manner other than as directed on the labeling (extralabel use) is not permitted.

➔
Residue Warning: At 200 g/ton use level or higher, withdraw 5 days before slaughter. Zero-day withdrawal period for lower use levels. Do not administer to turkeys producing eggs for human consumption.
➔

Combination Use:

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

VFD Issue Date: _____

VFD Expiration Date: _____

Veterinarian's Signature: _____

month/day/year

(Not to exceed 6 months from issuance date)