

Stafac[®] 50

Type A Medicated Article

Description:

Stafac Type A Medicated Article is an anti-infective for use in poultry and swine feeds to control enteric diseases. **Stafac** Type A Medicated Article has no withdrawal requirement.

Active Ingredient:

Virginiamycin11%
(Contains 50 g virginiamycin activity per lb)

Caution:

Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Mixing Directions:

Stafac 50 Type A Medicated Article must be thoroughly mixed in the feed to assure even distribution. Determine the appropriate amount of **Stafac** 50 Type A Medicated Article required from the table below. Dilute **Stafac** 50 Type A Medicated Article with a portion of one of the feed ingredients (enough to make a quantity equal to 10 lb per ton) before final mixing.

Safety:

Not for use in Laying Chickens.
Not for use in breeding swine over 120 lbs.
MUST be diluted in feed before use.
Feed continuously as sole ration.
Store at or below 25°C (77°F), Excursions Permitted Up To 40°C (104 °F).
Close container after use.

Registered Claims and Directions for Use:

| | Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed) | Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed) | Amount of virginiamycin activity provided to animal per lb of body weight per day |
|--|---|--|---|
| SWINE | | | |
| Treatment of swine dysentery in nonbreeding swine weighing over 120 lb | 100 g for 2 weeks | 2 lb | 2.27 mg |
| Treatment and control of swine dysentery in swine weighing up to 120 lb | 100 g for 2 weeks followed by 50 g | 2 lb 1 lb | 2.27 mg 1.14 mg |
| Aid in control of swine dysentery in swine weighing up to 120 lb For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred | 25 g | 8 oz. | 0.57 mg |
| CAUTION: Not for use in breeding swine over 120 lbs. | | | |
| POULTRY | | | |
| For use in broiler chickens Prevention of necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to virginiamycin | 20 g | 180 g | |
| Not for Use in Laying Chickens | | | |

Not for Human Use

Restricted Drug (California) - Use Only as Directed

HEALTHY ANIMALS. HEALTHY FOOD. HEALTHY WORLD.®

Phibro
ANIMAL HEALTH CORPORATION