



**FOR ANIMAL TREATMENT ONLY**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**PHIBROMONENSIN 400\***  
**400 G/ KG MONENSIN SODIUM**

PhibroMonensin is a technical active for addition to feed supplements and medicated premixes by formulators in order to achieve registered levels of monensin in the feed.

Indications

<b>Dairy Cattle</b>	For increased milk protein production As an aid in the control of ketosis As an aid in the prevention and control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>
<b>Beef Cattle</b>	As an aid in the prevention and control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>
<b>Replacement Heifers</b>	As an aid in the prevention and control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i>
<b>Sheep</b>	As an aid in the prevention and control of coccidiosis caused by, <i>E. ahsata</i> , <i>E. parva</i> , <i>E. ninakohlyakimovae</i> , <i>E. faurei</i> , <i>E. intricata</i> , <i>E. pallida</i> .
<b>Goats</b>	As an aid in the prevention and control of coccidiosis caused by <i>E. arloingi</i> .
<b>Broilers &amp; Layers</b>	As an aid in the prevention of coccidiosis in broilers and layer replacements caused by <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. maxima</i> , <i>E. mivati</i> , <i>E. necatrix</i> and <i>E. tenella</i> .
<b>Turkeys</b>	As an aid in the prevention of coccidiosis in growing turkeys only caused by <i>E. gallopavonis</i> , <i>E. adenoides</i> and <i>E. meleagrimitis</i> .

**25 Kg**

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

**DIRECTIONS FOR USE**

**This product must not be used for growth promotion in ruminants intended for human consumption.**

**Dairy Cattle:**

As an aid in the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuerni* and for increased milk protein production.

Complete Feeds: Thoroughly mix PhibroMonensin 400 into the ration to provide 300mg monensin per animal per day. The inclusion rate should be adjusted to take into account feed intake.

Supplements: Thoroughly mix PhibroMonensin 400 into the supplement to provide 100 - 300mg/kg per animal per day. The inclusion rate should be adjusted to take into account feed intake.

Ensure a minimum of 100 mg Monensin per animal per day is administered.

**Calves:**

Thoroughly mix PhibroMonensin 400 into the supplement to provide 0.5 -1.0 mg/kg live weight Monensin in the final feed. The inclusion rate should be adjusted to take into account feed intake.

Ensure a minimum of 100 mg Monensin per animal per day is administered.

**Beef Cattle and Replacement Heifers:**

For increased feed efficiency and /or rate of weight gain.

Complete Feeds: Thoroughly mix PhibroMonensin 400 into the ration to provide from 11 to 33 ppm monensin in the ration (90% dry matter).

Ensure a minimum dose of 100 mg Monensin per animal per day is administered.

Supplements: Thoroughly mix PhibroMonensin 400 into the ration to provide from 100 to 300 mg monensin per animal per day.

**Sheep and Goats:**

Complete Feeds: Thoroughly mix PhibroMonensin 400 into the ration to provide from 5 to 20 ppm monensin in the ration (90% dry matter).

Supplements: Thoroughly mix PhibroMonensin 400 into the ration to provide from 7.7 to 40 mg monensin per animal per day.

**Broiler and Layer Replacement Chickens:**

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella*

Thoroughly mix PhibroMonensin 400 into the ration to provide an equivalent of 100 to 120 g of monensin per tonne of complete feed.

**Growing Turkeys;**

A handwritten signature in blue ink, appearing to be 'John G. ...', located below the red stamp.

As an aid in the prevention of coccidiosis caused by *Eimeria adenoides*, *E.meleagrimitis* and *E. gallopavonis*.

Thoroughly mix PhibroMonensin 400 into the ration to provide an equivalent of 60 to 100 g of monensin per tonne of complete feed.

**Precautions: When mixing and handling PhibroMonensin 400 use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling.**

**Contraindications:**

**Do not treat cattle with products containing erythromycin, tiamulin or oleandomycin while using PhibroMonensin 400 Do not treat poultry with products containing tiamulin or oleandomycin while feeding PhibroMonensin 400. Severe growth depression may occur.**

**Caution:**

**Do not allow dogs, horses or other equines access to feed containing monensin. Consumption of feed containing monensin may be fatal to these animals.**

**Withholding Periods**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels of Agricultural Compounds.**

**Meat nil**

**Cattle Milk: nil. Sheep and goat milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.**

**Eggs: Eggs from treated birds must not be sold for human consumption for 10 days following the cessation of last treatment.**

**SAFETY DIRECTIONS:**

Poisonous if swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into feed wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves, goggles and half face piece respirator with dust cartridge or canister.[For mixing into daily supplements for individual animals when opening the container and mixing into feed wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves] After use and before eating, drinking or smoking, wash hands, arms and face



A handwritten signature in blue ink, appearing to be 'Timothy G. ...', located at the bottom right of the page.

thoroughly with soap and water. After each day's use, wash gloves, contaminated clothing, goggles and respirator and if rubber wash with detergent and warm water.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 03 4747000. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor

See Safety Data Sheet (SDS) for additional information.

**DISPOSAL:** Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burnt.

Registered Pursuant to the ACVM Act 1997 No A9493

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions  
APVMA Approval Number 55527

New Zealand Agent: David Marks  
PO Box 104095, Lincoln North, Auckland 0654

Registered to Phibro Animal Health Pty Ltd.  
Unit 1, 7-10 Denoci Close  
Wetherill Park, NSW 2164  
Australia

Store below 25°C (Air Conditioning). Protect from moisture.

Batch Number:

Expiry Date:



A handwritten signature in blue ink, appearing to be "David Marks", written over the bottom right portion of the red stamp.