



Nicarb® 25%

Type A Medicated Article with Microtracer®

Description:

Nicarb is a highly effective chemical anti-coccidial for the prevention of coccidiosis in broiler chickens. Preferred by customers for use in starter feeds, **Nicarb** prevents damaging cecal and intestinal species of *Eimeria* and results in improved gut health.

Active Ingredient:

Nicarbazin 25%

Registered Claims:

As an aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix*, and *E. brunetti*) coccidiosis in chickens.

Directions for Use:

Use nicarbazin Type C medicated feed as the only ration from the time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard.

Mixing Directions:

Nicarb (nicarbazin) **25%** should be thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for feed. Type C medicated feeds should contain 0.01 - 0.02% nicarbazin. Uniformly mix 0.8 - 1.6 pounds of **Nicarb 25%** with one of the feed ingredients to make at least 10 pounds of premix before final mixing. Mix the resulting premix with feed to make 1 ton (2000 lb) of finished feed containing 0.01 - 0.02% nicarbazin.

Warning:

Discontinue medication 4 days before marketing the birds for human consumption when used at levels less than or equal to 0.0125% and 5 days before, when used at levels greater than 0.0125% to allow for elimination of the drug from the edible tissue.

Contraindications:

Do not use in flushing mashers. Do not use as a treatment for outbreaks of coccidiosis.

Precautions

If losses exceed 0.5 percent in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist. Broilers fed feed medicated with **Nicarb 25%** may show reduced heat tolerance when exposed to high temperature and high humidity to which they have not been accustomed and under severe conditions, fatalities may result. An ample supply of drinking water and adequate ventilation will improve the birds' tolerance of heat.

Caution:

Do not feed to laying hens in production.